



Medications to Avoid Before Surgery or Injectables

We ask you stop taking the following medications and/or supplements with your physician's permission two weeks prior to surgery or injectable treatments (Botox, Xeomin, Juvederm, Kybella, Radiesse, Voluma, Volbella and Vollure) because these medications may promote bruising and/or bleeding. Please make a note of this prior to each appointment you make for an injectable, so we're able to meet your treatment expectations. If you are taking Coumadin, please notify us immediately and ONLY take acetaminophen (Tylenol) for pain prior to having your procedure.

ASPIRIN products include:

| | |
|--------------|-----------------|
| Alka-Seltzer | Anacin |
| Bayer | BC tablets |
| Cheracol | Cope |
| Coricidin | Davron compound |
| Dristan | Ecotrin |
| Empirin | Excedrin |
| Florinal | Midol |
| Percodan | Sine-Aid |
| Sine-Off | Soma compound |
| Stendin | Triaminicin |
| Vanquish | |

IBUPROFEN medications include:

| | |
|----------|--------|
| Advil | Alleve |
| Medipren | Motrin |
| Nuprin | Rufen |

ANITARTHRITIC medications include:

| | |
|----------|----------|
| Anaprox | Ansaid |
| Clinoril | Daypro |
| Dolobid | Feldene |
| Indocin | Naprosyn |
| Orudis | Relafen |
| Tolectin | Voltarin |

DIETARY SUPPLEMENTS include:

| | |
|-----------------------|----------------|
| Echinacea | Garlic tablets |
| Ginko biloba | Ginseng |
| St. John's Wort | Vitamin E |
| Omega 3/Cod liver oil | Red Wine |
| Weight loss products | |

**We encourage you to take 2,000 to 3,000mg of Vitamin C every day before your surgery and during recovery to alleviate bruising and swelling.*