



## Medications to Avoid Before Surgery or Injectables

*We ask you stop taking the following medications and/or supplements with your physician's permission two weeks prior to surgery or injectable treatments (Botox, Dysport, Redensity, Kysse, Juvederm, Voluma, Volux and Vollure) because these medications may promote bruising and/or bleeding. Please make a note of this prior to each appointment you make for an injectable, so we're able to meet your treatment expectations. If you are taking Coumadin, please notify us immediately and ONLY take acetaminophen (Tylenol) for pain prior to having your procedure.*

### **ASPIRIN products include:**

Alka-Seltzer	Anacin
Bayer	BC tablets
Cheracol	Cope
Coricidin	Davron compound
Dristan	Ecotrin
Empirin	Excedrin
Florinal	Midol
Percodan	Sine-Aid
Sine-Off	Soma compound
Stendin	Triaminicin
Vanquish	

### **IBUPROFEN medications include:**

Advil	Alleve
Medipren	Motrin
Nuprin	Rufen

### **WEIGHT LOSS medications include:**

Phentermine  
Tirzepatide (Mounjaro)  
Semaglutide (Ozempic/Wegovy)  
Liraglutide (Saxenda)

### **ANITARTHRITIC medications include:**

Anaprox	Ansaid
Clinoril	Daypro
Dolobid	Feldene
Indocin	Naprosyn
Orudis	Relafen
Tolectin	Voltarin

### **DIETARY SUPPLEMENTS include:**

Echinacea  
Ginko biloba  
St. John's Wort  
Omega 3/Cod liver oil  
Weight loss products  
Garlic tablets  
Ginseng  
Vitamin E  
Red Wine

*\*We encourage you to take 2,000 to 3,000mg of Vitamin C every day before your surgery and during recovery to alleviate bruising and swelling.*